arXiv:math/0510557v1 [math.DS] 26 Oct 2005

Some Inequalities satisfied by Periodical Solutions of Multi-Time Hamilton Equations

Iulian Duca, Constantin Udrişte

Abstract

The objective of this paper is to find some inequalities satisfied by periodical solutions of multi-time Hamilton systems, when the Hamiltonian is convex. To our knowledge, this subject of first-order field theory is still open.

Section 1 recall well-known facts regarding the equivalence between Euler-Lagrange equations and Hamilton equations and analyses the action that produces multi-time Hamilton equations, emphasizing the role of the polysymplectic structure. Section 2 extends two inequalities of [21] from a cube to parallelipiped and proves two inequalityes concerning multiple periodical solutions of multi-time Hamilton equations.

Key words: multi-time Hamilton action, Wirtinger inequality, convex Hamiltonian.

2000 Mathematics Subject Classification: 49S05, 70H05.

1 Multi-time Hamilton equations and polysymplectic structure

The paper studies the solutions with multiple periodicity of the Hamilton multi-time equations.

A function $u = (u^1, ..., u^n)$ with many variables $(t^1, ..., t^p)$, is multiple periodical with the period $T = (T^1, ..., T^p) \in R^p$ if

$$u\left(t^1 + k_1 T^1, ..., t^p + k_p T^p\right) = u\left(t^1, ..., t^p\right),$$

where $k_1, ..., k_p$ are integers. We consider the function u defined on the parallelepiped $T_0 = [0, T^1] \times [0, T^2] \times ... \times [0, T^p] \subset R^p$, with values in R^n. We will denote by $T = (T^1, ..., T^p) \in R^p$. The existence of the weak gradient of the function u assures the multiple periodicity of the function u. We use the Hilbert space H_T^1 attached to the Sobolev space $W_T^{1,2}$ of the functions $u \in L^2(T_0, R^n)$ which have a weak gradient $\dfrac{\partial u}{\partial t} \in L^2(T_0, R^n)$. The Wirtinger inequality from this paper has a specific form because of the multidimensional character of the definition domain T_0. The inequalities from theorems 3 and 4 constitute generalizations of some theorems of [5], from the particular case $p = 1$ to an arbitrary p.

The Euclidean structure on R^n is based on the scalar product $(u, v) = \delta_{ij} u^i v^j$, and the norm $|u| = \sqrt{\delta_{ij} u^i u^j}$. The Hilbert space H_T^1 is endowed with the scalar product

$$\langle u, v \rangle = \int_{T_0} \left(\delta_{ij} u^i(t) v^j(t) + \delta_{ij} \delta^{\alpha\beta} \frac{\partial u^i}{\partial t^\alpha}(t) \frac{\partial v^j}{\partial t^\beta}(t) \right) dt^1 \wedge ... \wedge dt^p,$$

and the corresponding norm $\sqrt{\langle u, u \rangle} = \|u\|$.

1.1 Multi-time Hamilton equations

We consider the multi-time variable $t = (t^1, ..., t^p) \in T_0 \subset R^p$, the functions $x^i : R^p \to R$, $(t^1, ..., t^p) \to x^i(t^1, ..., t^p)$, $i = 1, ..., n$, and the partial velocities $x_\alpha^i = \dfrac{\partial x^i}{\partial t^\alpha}$, $\alpha = 1, ..., p$.

Definition 1 The PDEs

$$\frac{\partial}{\partial t^\alpha} \frac{\partial L}{\partial x_\alpha^i} = \frac{\partial L}{\partial x^i}, \quad i = 1, ..., n, \quad \alpha = 1, ..., p$$

(second order PDEs system on the n-dimensional space) are called Euler-Lagrange equations for the Lagrangian

$$L : R^{p+n+np} \to R, \quad \left(t^\alpha, x^i, x_\alpha^i\right) \to L\left(t^\alpha, x^i, x_\alpha^i\right)$$

The Hamilton equations in the multi-time case are obtained using the partial derivatives (polymomenta)

$$p_k^\alpha = \frac{\partial L}{\partial x_\alpha^k} \tag{1}$$

and the Hamiltonian $H = p_k^\alpha x_\alpha^k - L$. If L satisfies some regularity conditions, then the system (1) defines a C^1 bijective transformation $x_\alpha^i \to p_i^\alpha$, called the Legendre transformation for the multi-time case. By this transformation we have

$$\frac{\partial H}{\partial p_i^\alpha} = x_\alpha^i + p_k^\beta \frac{\partial x_\beta^k}{\partial p_i^\alpha} - \frac{\partial L}{\partial x_\beta^k}\frac{\partial x_\beta^k}{\partial p_i^\alpha} = x_\alpha^i$$

$$\frac{\partial H}{\partial x^i} = p_k^\alpha \frac{\partial x_\alpha^k}{\partial x^i} - \frac{\partial L}{\partial x^i} - \frac{\partial L}{\partial x_\alpha^k}\frac{\partial x_\alpha^k}{\partial x^i} = -\frac{\partial L}{\partial x^i}.$$

Consequently, the $np + n$ Hamilton equations

$$\frac{\partial x^i}{\partial t^\alpha} = \frac{\partial H}{\partial p_i^\alpha},$$

$$\frac{\partial p_i^\alpha}{\partial t^\alpha} = -\frac{\partial H}{\partial x^i}$$

(summation after α), $i = 1, ..., n$, $\alpha = 1, ..., p$ are first order PDEs on the space R^{n+pn}, equivalent to the Euler-Lagrange equations on R^n.

There are different point of views to study these equations which appear in first-order field theory (see [1]-[3], [8]-[10], [12]-[22]). In our context, we need of Hilbert-Sobolev space methods for PDEs ([4], [6], [11]).

Let us write the multi-time Hamilton equations in the form

$$\delta_\beta^\alpha \delta_j^i \frac{\partial p_i^\beta}{\partial t^\alpha} + \frac{\partial H}{\partial x^j} = 0, \quad -\delta_\beta^\alpha \delta_j^i \frac{\partial x^j}{\partial t^\alpha} + \frac{\partial H}{\partial p_i^\beta} = 0, i, j = 1, ..., n; \alpha, \beta = 1, ..., p$$

or

$$(\delta \otimes J)\, \frac{\partial u}{\partial t} = -\nabla H, \tag{2}$$

where

$$\delta \otimes J = \begin{pmatrix} 0 & \delta_\beta^\alpha \delta_j^i \\ -\delta_\beta^\alpha \delta_j^i & 0 \end{pmatrix}, \frac{\partial u}{\partial t} = \begin{pmatrix} \dfrac{\partial x^j}{\partial t^\alpha} \\ \dfrac{\partial p_i^\beta}{\partial t^\alpha} \end{pmatrix}, \nabla H = \begin{pmatrix} \dfrac{\partial H}{\partial x^j} \\ \dfrac{\partial H}{\partial p_j^\beta} \end{pmatrix}.$$

The operator $\delta \otimes J$ is a polysymplectic structure acting on R^{np+np^2} with values in R^{n+np}. The (1,2)-block $\delta_\beta^\alpha \delta_j^i$ acts linearly by δ_j^i and tracely by δ_β^α.

The (2,1)-block $-\delta_\beta^\alpha \delta_j^i$ acts lineraly both by δ_j^i and δ_β^α. The operator $\delta \otimes J$ induces a multisymplectic PDE operator $(\delta \otimes J)\dfrac{\partial}{\partial t}$ which work as follows

$$(\delta \otimes J)\frac{\partial}{\partial t}\begin{pmatrix} x \\ p \end{pmatrix} : \begin{pmatrix} 0 & \delta_\beta^\alpha \delta_j^i \\ -\delta_\beta^\alpha \delta_j^i & 0 \end{pmatrix} \frac{\partial}{\partial t^\alpha}\begin{pmatrix} x^j \\ p_i^\beta \end{pmatrix} = \begin{pmatrix} \dfrac{\partial p_j^\alpha}{\partial t^\alpha} \\ -\dfrac{\partial x^j}{\partial t^\beta} \end{pmatrix}.$$

Repeating we obtain the square

$$(\delta \otimes J)\frac{\partial}{\partial t}\begin{pmatrix} \mathrm{div} p \\ -\dfrac{\partial x}{\partial t} \end{pmatrix} : \begin{pmatrix} 0 & \delta_\beta^\alpha \delta_j^i \\ -\delta_\beta^\alpha \delta_j^i & 0 \end{pmatrix} \begin{pmatrix} \dfrac{\partial^2 p_i^\gamma}{\partial t^\alpha \partial t^\gamma} \\ -\dfrac{\partial^2 x^j}{\partial t^\alpha \partial t\beta} \end{pmatrix} = \begin{pmatrix} -\Delta x^i \\ -\dfrac{\partial^2 p_i^\gamma}{\partial t^\beta \partial t^\gamma} \end{pmatrix}.$$

1.2 The action that produces multi-time Hamilton equations

We consider a Hamiltonian $H : T_0 \times R^n \times R^{np} \to R, (t,u) \to H(t,u)$ whose restriction $H(t,\cdot)$ is C^1 and convex.

Theorem 1 [21] *Let $u = (x,p)$. The action Ψ, whose Euler-Lagrange equations are the Hamilton equations, is*

$$\Psi(u) = \int_{T_0} \mathcal{L}\left(t, u, \frac{\partial u}{\partial t}\right) dt^1 \wedge ... \wedge dt^p,$$

$$\mathcal{L}\left(t, u, \frac{\partial u}{\partial t}\right) == -\frac{1}{2} G\left(\delta \otimes J\frac{\partial u}{\partial t}, u\right) - H(t,u),$$

where the scalar product is represented by the matrix

$$G = \begin{pmatrix} \delta_{ij} & 0 \\ 0 & \delta^{\beta\alpha}\delta_{ij} \end{pmatrix}$$

(standard Riemannian metric from R^{n+np}).

Proof. Indeed, the Euler-Lagrange equations produced by

$$\mathcal{L} = -\frac{1}{2}\left(\frac{\partial p_i^\alpha}{\partial t^\alpha} x^i - \frac{\partial x^i}{\partial t^\alpha} p_i^\alpha\right) - H(t,x,p) =$$

$$= -\frac{1}{2}\left(\frac{\partial p_i^\alpha}{\partial t^\alpha}, -\frac{\partial x^j}{\partial t^\beta}\right)\begin{pmatrix} \delta_{ij} & o \\ o & \delta^{\alpha\beta}\delta_{ij} \end{pmatrix}\begin{pmatrix} x^i \\ p_i^\alpha \end{pmatrix} - H\left(t, x\left(t\right), p\left(t\right)\right)$$

can be rewritten

$$\frac{1}{2}\frac{\partial p_i^\alpha}{\partial t^\alpha} = -\frac{1}{2}\frac{\partial p_i^\alpha}{\partial t^\alpha} - \frac{\partial H}{\partial x^i}\text{ , i.e. , }\frac{\partial p_i^\alpha}{\partial t^\alpha} = -\frac{\partial H}{\partial x^i}$$

and

$$\frac{\partial x^i}{\partial t^\alpha} = \frac{\partial H}{\partial p_i^\alpha}.$$

2 Basic inequalities

2.1 Wirtinger multi-time inequality

In $L^2\left(T_0, R^n\right)$ we use the scalar product

$$\langle u, v\rangle = \int_{T_0}\left(\delta_{ij}u^i v^j\right)dt^1 \wedge ... \wedge dt^p$$

and the norm $\|u\|_{L^2} = \sqrt{\langle u, u\rangle}$. Similarly, in $L^2\left(T_0, C^n\right)$ we use the scalar product

$$\langle u, v\rangle = \int_{T_0}\left(\delta_{ij}u^i \overline{v}^j\right)dt^1 \wedge ... \wedge dt^p$$

and the norm $\|u\|_{L^2} = \sqrt{\langle u, \overline{u}\rangle}$.

Let us extend the Theorem 4.4 from [21] to the parallelipiped T_0.

Theorem 2 *Any function u from H_T^1 with mean zero satisfies the inequality*

$$\int_{T_0}\left|u\left(t\right)\right|^2 dt^1 \wedge ... \wedge dt^p \leq \frac{\left(\max\limits_i\left\{T^i\right\}\right)^2}{4\pi^2}\int_{T_0}\left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p.$$

Proof. We express the function u as the sum of a multiple Fourier series

$$u\left(t\right) = \left(u^1\left(t^1, ..., t^p\right), ..., u^n\left(t^1, ..., t^p\right)\right)$$

$$= \left(\sum_{j_1^1,...,j_1^p \in Z^p} C^1_{j_1^1,...,j_1^p} e^{i\left(\frac{2\pi}{T^1}\cdot j_1^1 t^1 + ... + \frac{2\pi}{T^p}\cdot j_1^p t^p\right)}, \right.$$

$$\left. ..., \sum_{j_n^1,...,j_n^p \in Z^p} C^n_{j_n^1,...,j_n^p} e^{i\left(\frac{2\pi}{T^1}\cdot j_n^1 t^1 + ... + \frac{2\pi}{T^p}\cdot j_n^p t^p\right)} \right).$$

We calculate the square of the norm

$$(u,u) = \int_{T_0} \left(u\left(t\right), \overline{u\left(t\right)}\right) dt^1 \wedge ... \wedge dt^p =$$

$$= \sum_{j_1^1,...,j_1^p \in Z^p} \left(C^1_{j_1^1,...,j_1^p}\right)^2 \int_0^{T^1} e^{i\frac{2\pi}{T^1}\left(j_1^1 - j_1^1\right)t^1} dt^1 ... \int_0^{T^p} e^{i\frac{2\pi}{T^p}\left(j_1^p - j_1^{p1}\right)t^p} dt^p + ...$$

$$... + \sum_{j_n^1,...,j_n^p \in Z^p} \left(C^1_{j_n^1,...,j_n^p}\right)^2 \int_0^{T^1} e^{i\frac{2\pi}{T^1}\left(j_n^1 - j_n^1\right)t^1} dt^1 ... \int_0^{T^p} e^{i\frac{2\pi}{T^p}\left(j_n^p - j_n^p\right)t^p} dt^p$$

$$= \sum_{j_1^1,...,j_1^p \in Z^p} \left(C^1_{j_1^1,...,j_1^p}\right)^2 T^1 ... T^p + ... + \sum_{j_1^1,...,j_1^p \in Z^p} \left(C^n_{j_n^1,...,j_n^p}\right)^2 T^1 ... T^p.$$

If we denote

$$C_{k_1,...,k_p} = \left(C^1_{k_1,...,k_p}, ..., C^n_{k_1,...,k_p}\right)$$

and

$$u = \sum_{(k_1,...,k_p) \in Z^p} C_{k_1,...,k_p} e^{i2\pi\left(\frac{k_1}{T^1}t_1 + ... + \frac{k_p}{T^p}t_p\right)},$$

we find

$$(u,u) = \sum_{(k_1,...,k_p) \in Z^p} \left|C_{k_1,...,k_p}\right|^2 T^1 ... T^p.$$

Similarly, we consider the scalar product

$$\left(\frac{\partial u}{\partial t}, \frac{\partial v}{\partial t}\right) = \int_{T_0} \delta_{ij}\delta^{\alpha\beta} \frac{\partial u^i}{\partial t^\alpha} \overline{\frac{\partial v^j}{\partial t^\beta}} dt^1 \wedge ... \wedge dt^p.$$

It follows the square of the norm

$$\left(\frac{\partial u}{\partial t},\frac{\partial u}{\partial t}\right)$$

$$=\sum_{(k_1,...,k_p)\in Z^p}T^1...T^p\left[\left(C^1_{k_1...k_p}\right)^2\left(\frac{2\pi k_1}{T^1}\right)^2+...+\left(C^1_{k_1...k_p}\right)\left(\frac{2\pi k_p}{T^p}\right)^2+...\right.$$

$$\left....+\left(C^n_{k_1...k_p}\right)^2\left(\frac{2\pi k_1}{T^1}\right)^2+...+\left(C^n_{k_1...k_p}\right)^2\left(\frac{2\pi k_p}{T^p}\right)^2\right]$$

$$=\sum_{(k_1,...,k_p)\in Z^p}T^1...T^p\left[\left|C_{k_1...k_p}\right|^2\left(\frac{2\pi k_1}{T^1}\right)^2+...+\left|C_{k_1...k_p}\right|^2\left(\frac{2\pi k_p}{T^p}\right)^2\right]$$

$$=\sum_{(k_1,...,k_p)\in Z^p}T^1...T^p\left[\left|C_{k_1...k_p}\right|^2 4\pi^2\left(\left(\frac{k_1}{T^1}\right)^2+...+\left(\frac{k_p}{T^p}\right)^2\right)\right]$$

$$\geq\frac{4\pi^2}{\left(\max\limits_i\left\{T^i\right\}\right)^2}\sum_{(k_1,...,k_p)\in Z^p}T^1...T^p\left|C_{k_1...k_p}\right|^2\left(k_1^2+...+k_1^2\right)$$

$$\geq\frac{4\pi^2}{\left(\max\limits_i\left\{T^i\right\}\right)^2}\sum_{(k_1,...,k_p)\in Z^p}T^1...T^p\left|C_{k_1...k_p}\right|^2$$

$$\geq\frac{4\pi^2}{\left(\max\limits_i\left\{T^i\right\}\right)^2}\int_{T_0}\left|u\left(t\right)\right|^2dt^1\wedge...\wedge dt^p.$$

Consequently

$$\int_{T_0}\left|u\left(t\right)\right|^2dt^1\wedge...\wedge dt^p\leq\frac{\left(\max\limits_i\left\{T^i\right\}\right)^2}{4\pi^2}\int_{T_0}\left|\frac{\partial u}{\partial t}\right|^2dt^1\wedge...\wedge dt^p,$$

and this ends the proof.

2.2 An estimate of the quadratic form $\int_{T_0}\left(\delta\otimes J\frac{\partial u}{\partial t},u\left(t\right)\right)dt^1\wedge...\wedge dt^p$

Let us extend the Theorem 4.5 from [21] to the parallelipiped T_0.

Theorem 3 *For any $u\in H_T^1$ we have*

$$\int_{T_0}\left(\delta\otimes J\frac{\partial u}{\partial t},u\left(t\right)\right)dt^1\wedge...\wedge dt^p$$

$$\geq-\frac{\sqrt{p}\max\limits_i\left\{T^i\right\}}{2\pi}\int_{T_0}\left|\frac{\partial u}{\partial t}\right|^2dt^1\wedge...\wedge dt^p.$$

Proof. We denote $\widetilde{u}\left(t\right)=u\left(t\right)-\int_{T_0}u\left(t\right)dt^1\wedge...\wedge dt^p$.. By using the Cauchy-Schwarz inequality and the multiple periodicity of u we obtain the inequality

$$\int_{T_0}\left(\delta\otimes J\frac{\partial u}{\partial t},u\left(t\right)\right)dt^1\wedge...\wedge dt^p$$

$$=\int_{T_0}\left(\delta\otimes J\frac{\partial u}{\partial t},\widetilde{u}\left(t\right)+\int_{T_0}u\left(t\right)dt^1\wedge...\wedge dt^p\right)dt^1\wedge...\wedge dt^p$$

$$=\int_{T_0}\left(\delta\otimes J\frac{\partial u}{\partial t},\widetilde{u}\left(t\right)\right)dt^1\wedge...\wedge dt^p$$

$$+\int_{T_0}\left(\delta\otimes J\frac{\partial u}{\partial t},\int_{T_0}u\left(t\right)dt^1\wedge...\wedge dt^p\right)dt^1\wedge...\wedge dt^p$$

$$=\int_{T_0}\left(\delta\otimes J\frac{\partial u}{\partial t},\widetilde{u}\left(t\right)\right)dt^1\wedge...\wedge dt^p$$

$$\geq-\left(\int_{T_0}\left|\delta\otimes J\frac{\partial u}{\partial t}\right|^2dt^1\wedge...\wedge dt^p\right)^{\frac{1}{2}}\left(\int_{T_0}\left|\widetilde{u}\left(t\right)\right|^2dt^1\wedge...\wedge dt^p\right)^{\frac{1}{2}}$$

From the inequality given by the Theorem 2 we have

$$\int_{T_0} |\widetilde{u}(t)|^2 \, dt^1 \wedge ... \wedge dt^p \leq \frac{\left(\max\limits_i \left\{T^i\right\}\right)^2}{4\pi^2} \int_{T_0} \left|\frac{\partial \widetilde{u}}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p$$

$$= \frac{\left(\max\limits_i \left\{T^i\right\}\right)^2}{4\pi^2} \int_{T_0} \left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p.$$

Because $\left|\delta \otimes J\frac{\partial u}{\partial t}\right|^2 \leq p\left|\frac{\partial u}{\partial t}\right|^2$, we obtain

$$\int_{T_0} \left(\delta \otimes J\frac{\partial u}{\partial t}, u(t)\right) dt^1 \wedge ... \wedge dt^p$$

$$\geq -\left(\int_{T_0} \left|\delta \otimes J\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p\right)^{\frac{1}{2}} \frac{\max\limits_i \left\{T^i\right\}}{2\pi}$$

$$\cdot \left(\int_{T_0} \left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p\right)^{\frac{1}{2}}$$

$$\geq -\sqrt{p} \ \frac{\max\limits_i \left\{T^i\right\}}{2\pi} \int_{T_0} \left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p.$$

2.3 Inequalities satisfied by periodical solutions of multi-time Hamilton equations

Let us find properties of solutions of $(\delta \otimes J)\frac{\partial u}{\partial t} + \nabla H(t, u(t)) = 0$ a.e. on T_0 satisfying the boundary conditions

$$u\,|_{S^+} = u\,|_{S^-},$$

were S^+ and S^- are opposite sides of the parallelipiped T_0. Practically, we refer to bounds for such solutions.

Theorem 4 *We consider the Hamiltonian*

$$H : T_0 \times R^{n+np} \rightarrow R, (t,u) \rightarrow H(t,u)$$

like a measurable function in t for any $u \in R^{n+np}$, and C^1 *convex in u for any*

$$t \in T_0 = \left[0, T^1\right] \times ... \times [0, T^p] \subset R^p.$$

If there exists the constants

$$\alpha \in \left(0, \frac{\pi}{\sqrt{p} \max\limits_i \left\{T^i\right\}}\right), \beta \geq 0, \gamma \geq 0, \delta \geq 0$$

such that

$$\delta |u| + \beta \leq H(t,u) \leq \frac{\alpha}{2} |u|^2 + \gamma$$

for all $t \in T_0$ *and* $u \in R^{n+np}$, *then, any multiple periodical solution*

$$u = (x_i, p_i^\alpha), i = 1, ..., n, \alpha = 1, ..., p,$$

of the equation

$$\delta \otimes J \frac{\partial u}{\partial t} + \nabla H(t, u(t)) = 0, \tag{3}$$

verifies the inequalities

$$\int_{T_0} \left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p \leq \frac{2\alpha(\beta+\gamma)\pi T^1 ... T^p}{\pi - \alpha \max\limits_i \left\{T^i\right\} \sqrt{p}} \tag{4}$$

$$\int_{T_0} |u(t)| dt^1 \wedge ... \wedge dt^p \leq \frac{\pi T^1 ... T^p (\beta+\gamma)}{\delta\left(\pi - \alpha \max\limits_i \left\{T^i\right\} \sqrt{p}\right)}. \tag{5}$$

Proof. From the inequality

$$\delta |u|^2 - \beta \leq H(t,u) \leq \frac{\alpha}{2} |u|^2 + \gamma$$

we obtain

$$-\beta \leq H(t,u) \leq \alpha 2^{-1} |u|^2 + \gamma.$$

By applying [5, Proposition 2.2], considering $F(u) = H(t,u)$, $p = q = 2$, $v = \nabla H(t,u)$ we obtain

$$\frac{1}{2\alpha}\left|\nabla H(t,u)\right|^2 \leq \left(\nabla H(t,u), u\right) + \beta + \gamma.$$

Because u is the solution of the equation (3), we have $\nabla H(t,u) = -\delta \otimes J\dfrac{\partial u}{\partial t}$ and the previous inequality becomes

$$\frac{1}{2\alpha}\left|-\delta \otimes J\frac{\partial u}{\partial t}\right|^2 \leq \left(-\delta \otimes J\frac{\partial u}{\partial t}, u\right) + \beta + \gamma. \tag{6}$$

In the hypothesis' conditions, by integration of the inequality (6) we have

$$\frac{1}{2\alpha}\int_{T_0}\left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p$$

$$+\int_{T_0}\left(\delta \otimes J\frac{\partial u}{\partial t}, u\right) dt^1 \wedge ... \wedge dt^p \leq (\beta+\gamma)\, T^1 ... T^p.$$

By using the inequality from Theorem 3, we have

$$\frac{1}{2\alpha}\int_{T_0}\left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p$$

$$-\frac{\sqrt{p}\max_i\{T^i\}}{2\pi}\int_{T_0}\left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p \leq (\beta+\gamma)\, T^1 ... T^p.$$

So

$$\left(\frac{1}{2\alpha} - \frac{\sqrt{p}\max\limits_i\left\{T^i\right\}}{2\pi}\right)\int_{T_0}\left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p$$

$$\leq (\beta+\gamma)\, T^1 ... T^p$$

and, as consequence

$$\int_{T_0}\left|\frac{\partial u}{\partial t}\right|^2 dt^1 \wedge ... \wedge dt^p \leq \frac{2\pi\alpha(\beta+\gamma)\, T^1 ... T^p}{\pi - \alpha\max\limits_i\left\{T^i\right\}\sqrt{p}}.$$

By integration, the inequality

$$\delta |u| - \beta \leq H(t,u)$$

produces

$$\delta \int_{T_0} |u(t)| \, dt^1 \wedge ... \wedge dt^p - \beta T^1 ... T^p \leq \int_{T_0} H(t,u) \, dt^1 \wedge ... \wedge dt^p.$$

Because $H(t,u)$ is convex in u,

$$H(t,u) - H(t,0) \leq (\nabla H(t,u(t)), u(t)),$$

we obtain

$$\begin{aligned}
&\int_{T_0} H(t,u(t)) \, dt^1 \wedge ... \wedge dt^p \\
&\leq \int_{T_0} [H(t,0) + (\nabla H(t,u(t)), u(t))] \, dt^1 \wedge ... \wedge dt^p \\
&\leq \gamma T^1 ... T^p - \int_{T_0} \left(\delta \otimes J \frac{\partial u}{\partial t}, u(t) \right) dt^1 \wedge ... \wedge dt^p \\
&\leq \gamma T^1 ... T^p + \frac{\sqrt{p} \max_i \left\{ T^i \right\}}{2\pi} \int_{T_0} \left| \frac{\partial u}{\partial t} \right|^2 dt^1 \wedge ... \wedge dt^p \\
&\leq \gamma T^1 ... T^p + \frac{\sqrt{p} \max_i \left\{ T^i \right\} 2\pi\alpha (\beta + \gamma) T^1 ... T^p}{2\pi \left(\pi - \alpha \max_i \left\{ T^i \right\} \sqrt{p} \right)}.
\end{aligned}$$

By consequence

$$\begin{aligned}
&\int_{T_0} |u(t)| \, dt^1 \wedge ... \wedge dt^p \\
&\leq \frac{1}{\delta} \left(\beta T^1 ... T^p + \gamma T^1 ... T^p + \frac{\sqrt{p} \max_i \left\{ T^i \right\} \alpha (\beta + \gamma)}{\pi - \alpha \max_i \left\{ T^i \right\} \sqrt{p}} T^1 ... T^p \right),
\end{aligned}$$

meaning that

$$\int_{T_0} |u(t)| \, dt^1 \wedge ... \wedge dt^p \leq \frac{(\beta + \gamma) T^1 ... T^p \pi}{\delta \left(\pi - \alpha \max_i \left\{ T^i \right\} \sqrt{p} \right)}$$

and the proof ends.

References

[1] I. Duca, A-M. Teleman, C. Udrişte: *Poisson-Gradient Dynamical Systems with Convex Potential*, Proceedings of the 3-rd International Colloquium " Mathematics in Engineering and Numerical Physics ", 7-9 October, 2004, Bucharest.

[2] M. Forger, C. Paufler, H. Romer, *The Poisson bracket for Poisson forms in multisymplectic field theory*, Reviews in Mathematical Physics, 15, 7 (2003), 705-743.

[3] I. V. Kanatchikov: *Geometric (pre)quantization in the polysymplectic approach to field theory*, arXiv: hep-th/0112263 v3, 3 Jun 2002, 1-12; Differential Geometry and Its Applications, Proc. Conf., Opava (Czech Republic), August 27-31, 2001, Silesian University Opava, 2002.

[4] L. V. Kantorovici, G. P. Akilov: *Analiză funcţională*, Editura ştiinţifică şi enciclopedică, Bucureşti, 1980.

[5] J. Mawhin, M. Willem: *Critical Point Theory and Hamiltonian Systems*, Springer-Verlag, 1989.

[6] S. G. Mihlin: *Ecuaţii liniare cu derivate parţiale*, Editura ştiinţifică şi enciclopedică, Bucureşti, 1983.

[7] M. Neagu, C. Udrişte : *From PDE Systems and Metrics to Geometric Multi-time Field Theories*, Seminarul de Mecanică, Sisteme Dinamice Diferenţiale, 79, Universitatea de Vest din Timişoara, 2001.

[8] P. J. Olver: *The Equivalence Problem and Canonical Forms for Quadratic Lagrangians*, Advances in Applied Mathematics, 9, (1998), 226-227.

[9] C. Paufler, H. Romer: *De Donder-Weyl equations and multisymplectic geometry*, arXiv: math-ph/0506022 v2, 20 Jun 2005.

[10] N. Roman-Roy, *Multisymplectic Lagrangian and Hamiltonian formalism of first -order classical field theories*, arXiv: math-ph/0107019 v1, 20 Jul 2001, vol XX (XXXX), No. X, 1-9.

[11] R. E. Showalter: *Hilbert Space Methods for Partial Differential Equations*, Electronic Journal of Differential Equations Monograph 01, 1994.

[12] A.-M. Teleman, C. Udrişte: *On Hamiltonian Formalisms in Mathematical Physics*, 4-th International Conference of Balkan Society of Geometers, Aristotle University of Thessaloniki, 26-30 June, 2002.

[13] C. Udrişte: *Nonclassical Lagrangian Dynamics and Potential Maps*, Conference in Mathematics in Honour of Professor Radu Roşca on the occasion of his Ninetieth Birthday, Katholieke University Brussel, Katholieke University Leuwen, Belgium, December 11-16, 1999; http://xxx.lanl.gov/math.DS/0007060, (2000).

[14] C. Udrişte: *Geometric Dynamics*, Kluwer Academic Publishers, Dodrecht /Boston /London, Mathematics and Its Applications, 513, (2000); Southeast Asian Bulletin of Mathematics, Springer-Verlag 24 (2000).

[15] C. Udrişte: *Solutions of DEs and PDEs as Potential Maps Using First Order Lagrangians*, Centenial Vrânceanu, Romanian Academy, University of Bucharest, June 30-July 4, (2000); http://xxx.lanl.gov/math.DS/0007061, (2000); Balkan Journal of Geometry and Its Applications 6, 1, 93-108, 2001.

[16] C. Udrişte: *From integral manifolds and metrics to potential maps*, Atti del'Academia Peloritana dei Pericolanti,Classe 1 di Scienze Fis. Mat. e Nat., 81-82, A 01006 (2003-2004), 1-14.

[17] C. Udrişte, I. Duca: *Periodical Solutions of Multi-Time Hamilton Equations,* Analele Universitaţii Bucureşti, 55, 1 (2005), 179-188.

[18] C. Udrişte, I. Duca: *Poisson-gradient Dynamical Systems with Bounded Non-linearity*, manuscript.

[19] C. Udrişte, M. Neagu: *From PDE Systems and Metrics to Generalized Field Theories*, http://xxx.lanl.gov/abs/math.DG/0101207.

[20] C. Udrişte, M. Postolache: *Atlas of Magnetic Geometric Dynamics*, Geometry Balkan Press, Bucharest, 2001.

[21] C. Udrişte, A.-M. Teleman: *Hamilton Approaches of Fields Theory*, IJMMS, 57 (2004), 3045-3056; ICM Satelite Conference in Algebra and Related Topics, University of Hong-Kong , 13-18.08.02.

[21] C. Udrişte, M. Ferrara, D. Opriş, Economic Geometric Dynamics, Geometry Balkan Press, Bucharest, 2004.

University Politehnica of Bucharest
Departament of Mathematics
Splaiul Independenţei 313
060042 Bucharest, Romania
udriste@mathem.pub.ro